arXiv:1109.6476v1 [math.CA] 29 Sep 2011

Bifurcation of limit cycles by perturbing a piecewise linear Hamiltonian system with a homoclinic loop [*]

Liang Feng[a,b][†], Han Maoan[a,c] [‡]and Valery G. Romanovski[c,d][§]

[a]The Institute of Mathematics, Shanghai Normal University, Shanghai 200234, PR China

[b]The Institute of Mathematics, Anhui Normal University, Wuhu 241000, PR China

[c]Faculty of Natural Science and Mathematics,
University of Maribor, SI-2000 Maribor, Slovenia

[d]Center for Applied Mathematics and Theoretical Physics,
University of Maribor, SI-2000 Maribor, Slovenia

Abstract. In this paper, we study limit cycle bifurcations for a kind of non-smooth polynomial differential systems by perturbing a piecewise linear Hamiltonian system with a center at the origin and a homoclinic loop around the origin. By using the first Melnikov function of piecewise near-Hamiltonian systems, we give lower bounds of the maximal number of limit cycles in Hopf and homoclnic bifurcations, and derive an upper bound of the number of limit cycles that bifurcate from the periodic annulus between the center and the homoclinic loop up to the first order in ε. In the case when the degree of perturbing terms is low, we obtain a precise result on the number of zeros of the first Melnikov function.

MSC. 34C05; 34C07; 37G15

Keywords: Limit cycle; homoclinic loop; Melnikov function; bifurcation; piecewise smooth system.

1 Introduction and main results

There are many problems in mechanics, electrical engineering and the theory of automatic control which are described by non-smooth systems, see for instance the works of A.F.Filippov [1], Andronov et al [2], Kunze [3] and the references therein. Recently, a good deal of work has been done to study bifurcations in non-smooth systems including Hopf, homoclinic and subharmonic bifurcations, etc.. In [4-6] Hopf bifurcation for

---

[*] The project supported by the National Natural Science Foundation of China (10971139) and the Slovenian Research Agency.

[†] *E-mail address*: liangfeng741018@163.com

[‡] Author for correspondence. *E-mail address*: mahan@shnu.edu.cn.

[§] *E-mail address*: valery.romanovsky@uni-mb.si.

non-smooth systems was studied by developing new methods for computing Lyapunov constants. The Melnikov method for Hopf and homoclinic bifurcations was extended to non-smooth systems in [7-9]. The method of averaging has also been extended to non-smooth systems in [10]. However, so far there are few papers in the literature studying homoclinic bifurcations inside the class of piecewise polynomial differential systems. In this work, we study this problem by using the first order Melnikov function of piecewise near-Hamiltonian systems.

In [7], Liu and Han considered a general form of a piecewise near-Hamiltonian system on the plane

$$\left\{\begin{array}{l} \dot{x} = H_y + \varepsilon p(x, y, \delta), \\ \dot{y} = -H_x + \varepsilon q(x, y, \delta), \end{array}\right. \tag{1.1}$$

where

$$H(x, y) = \left\{\begin{array}{l} H^+(x, y),\ x \geq 0, \\ H^-(x, y),\ x < 0, \end{array}\right.$$

$$p(x, y, \delta) = \left\{\begin{array}{l} p^+(x, y, \delta),\ x \geq 0, \\ p^-(x, y, \delta),\ x < 0, \end{array}\right.$$

$$q(x, y, \delta) = \left\{\begin{array}{l} q^+(x, y, \delta),\ x \geq 0, \\ q^-(x, y, \delta),\ x < 0, \end{array}\right.$$

$H^\pm$, $p^\pm$ and $q^\pm$ are C^∞, $\varepsilon > 0$ is small, $\delta \in D \subset R^m$ is a vector parameter with D compact. This system has two subsystems

$$\left\{\begin{array}{l} \dot{x} = H_y^+ + \varepsilon p^+(x, y, \delta), \\ \dot{y} = -H_x^+ + \varepsilon q^+(x, y, \delta), \end{array}\right. \tag{1.1a}$$

and

$$\left\{\begin{array}{l} \dot{x} = H_y^- + \varepsilon p^-(x, y, \delta), \\ \dot{y} = -H_x^- + \varepsilon q^-(x, y, \delta), \end{array}\right. \tag{1.1b}$$

which are called the right subsystem and the left subsystem, respectively. Suppose that $(1.1)|_{\varepsilon=0}$ has a family of periodic orbits around the origin and satisfies the following two assumptions.

Assumption (I): There exist an interval $J = (\alpha, \beta)$, and two points $A(h) = (0, a(h))$ and $A_1(h) = (0, a_1(h))$ such that for $h \in J$

$$H^+(A(h)) = H^+(A_1(h)) = h,\ H^-(A(h)) = H^-(A_1(h)) = \widetilde{h},\ a(h) \neq a_1(h).$$

Assumption (II): The subsystem $(1.1\text{a})|_{\varepsilon=0}$ has an orbital arc L_h^+ starting from $A(h)$ and ending at $A_1(h)$ defined by $H^+(x, y) = h, x \geq 0$; the subsystem $(1.1\text{b})|_{\varepsilon=0}$ has an orbital arc L_h^- starting from $A_1(h)$ and ending at $A(h)$ defined by $H^-(x, y) = H^-(A_1(h)), x < 0$.

Under assumptions (I) and (II), $(1.1)|_{\varepsilon=0}$ has a family of non-smooth periodic orbits $L_h = L_h^+ \cup L_h^-$, $h \in J$. For definiteness, we assume the orbits L_h for $h \in J$ orientate anticlockwise. See Fig. 1.

By Theorem 1.1 in [7], the first order Melnikov function of system (1.1) has the form

$$M(h, \delta) = \frac{H_y^+(A)}{H_y^-(A)} \Big[\frac{H_y^-(A_1)}{H_y^+(A_1)} \int_{L_h^+} q^+ dx - p^+ dy + \int_{L_h^-} q^- dx - p^- dy \Big],\ h \in J. \tag{1.2}$$



Fig. 1. The closed orbits of $(1.1)|_{\varepsilon=0}$.

Also, we know from [7] that if $M(h,\delta)$ has at most k zeros in h on the interval (α,β) for all $\delta \in D$, then (1.1) has at most k limit cycles bifurcated from the open annulus $\bigcup_{\alpha<h<\beta} L_h$.

Let

$$\begin{aligned} &M^+(h,\delta) = \textstyle\int_{L_h^+} q^+dx - p^+dy,\ M^-(h,\delta) = \int_{L_h^-} q^-dx - p^-dy, \\ &\widetilde{M}^-(\widetilde{h},\delta) = \int_{H^-(x,y)=\widetilde{h},\ x\le 0} q^-dx - p^-dy, \end{aligned} \tag{1.3}$$

where $\widetilde{h}$ is given in assumption (I). Then we know

$$M^-(h,\delta) = \widetilde{M}^-(H^-(A_1(h)),\delta),\ h \in J. \tag{1.4}$$

As in the smooth case, a very important issue associated with (1.1) is to find the number of limit cycles and their distribution. In [7], Liu and Han consider a piecewise polynomial system of the form

$$(\dot{x},\dot{y}) = \begin{cases} (b^+y + \varepsilon p_n^+(x,y,\delta), -b^+x + \varepsilon q_n^+(x,y,\delta)),\ x \ge 0, \\ (b^-y + \varepsilon p_n^-(x,y,\delta), -b^-x + \varepsilon q_n^-(x,y,\delta)),\ x < 0, \end{cases} \tag{1.5}$$

where $b^\pm > 0$, $p_n^\pm$ and $q_n^\pm$ are arbitrary polynomials of degree n. It was proved that the maximal number of limit cycles of the above system is n up to the first order in ε. We know the unperturbed system of (1.5) has a global center at the origin. Then, an interesting question is how many limit cycles system (1.1) can have if $(1.1)|_{\varepsilon=0}$ is a linear piecewise system with a homoclinic loop and $p^\pm$ and $q^\pm$ are arbitrary polynomials with degree n.

In this paper, we take

$$H^+(x,y) = \frac{1}{2}\big((x-1)^2 - y^2\big),\ x \ge 0, \tag{1.6}$$

$$H^-(x,y) = -\frac{1}{2}(x^2+y^2),\ x < 0, \tag{1.7}$$

and suppose

$$p(x,y)=\begin{cases} p^+(x,y)=\sum\limits_{i+j=0}^{n} a_{ij}^+x^iy^j,\ x\geq 0,\\ p^-(x,y)=\sum\limits_{i+j=0}^{n} a_{ij}^-x^iy^j,\ x<0,\end{cases}$$
$$q(x,y)=\begin{cases} q^+(x,y)=\sum\limits_{i+j=0}^{n} b_{ij}^+x^iy^j,\ x\geq 0,\\ q^-(x,y)=\sum\limits_{i+j=0}^{n} b_{ij}^-x^iy^j,\ x<0.\end{cases} \tag{1.8}$$

Under (1.6) and (1.7), the system (1.1) becomes

$$\begin{cases} \dot{x}=-y+\varepsilon p^+(x,y),\\ \dot{y}=1-x+\varepsilon q^+(x,y),\end{cases} x\geq 0, \qquad \begin{cases} \dot{x}=-y+\varepsilon p^-(x,y),\\ \dot{y}=x+\varepsilon q^-(x,y),\end{cases} x<0. \tag{1.9}$$

For system $(1.9)|_{\varepsilon=0}$, there exist a family of periodic orbits as follows

$$L_h=\{(x,y)|H^+(x,y)=\frac{h}{2},\ x\geq 0\}\cup\{(x,y)|H^-(x,y)=\frac{\widetilde{h}}{2},\ x<0\},$$

with $\widetilde{h}=h-1$, $0<h<1$. For the sake of convenience, here we use $h/2$ instead of h. If $h\to 1$, L_h approaches the origin which is an elementary center of parabolic-focus type(see [4] for the definition). And if $h\to 0$, $L_h\to L_0$, where L_0 is a compound homoclinic loop with a saddle $S_1(1,0)$. See Fig. 2.



Fig. 2. Phase portrait of system $(1.9)|_{\varepsilon=0}$.

Note that $H_y^+(0,y)\equiv H_y^-(0,y)$ for $-1<y<1$. Then by (1.2), we have the first order Melnikov function of system (1.9) satisfying

$$M(\frac{h}{2})=\int_{\widehat{AA_1}} q^+dx-p^+dy+\int_{\widehat{A_1A}} q^-dx-p^-dy\equiv\bar{M}(h), \tag{1.10}$$

where $0 < h < 1$, and

$$\begin{gathered} A = (0, -\sqrt{1-h}),\ A_1 = (0, \sqrt{1-h}), \\ \widehat{AA_1} = \{(x,y)|H^+(x,y) = \tfrac{h}{2},\ x \geq 0\}, \\ \widehat{A_1A} = \{(x,y)|H^-(x,y) = \tfrac{h-1}{2},\ x < 0\}. \end{gathered}$$

Let $Z(n)$ denote the maximal number of zeros of the non-zero function $\bar{M}(h)$ on the open interval $(0,1)$ for all possible p and q satisfying (1.8), which is the maximal number of limit cycles of (1.9) bifurcated from the periodic annulus $\bigcup_{0<h<1} L_h$ for all possible p and q satisfying (1.8) when $\bar{M}(h)$ is not identically zero. Let $N_{Hopf}(n)$ and $N_{homoc}(n)$ denote, respectively, the maximal number of limit cycles bifurcated in Hopf bifurcation near the origin and in homoclinic bifurcation near L_0 for all possible p and q satisfying (1.8). Then our main results can be stated as follows.

Theorem 1.1. *For any* $n \geq 1$ *we have*

(1) $N_{Hopf}(n) \geq n + [\frac{n+1}{2}]$.

(2) $N_{homoc}(n) \geq n + [\frac{n+1}{2}]$.

Theorem 1.2. *For* $n = 1, 2, 3, 4,$ *we have* $Z(n) = n + [\frac{n+1}{2}]$.

Theorem 1.3. *For any* $n \geq 5$ *we have* $n + [\frac{n+1}{2}] \leq Z(n) \leq 2n + [\frac{n+1}{2}]$.

2 Preliminary

In this section we give an expression of the first order Melnikov function $\bar{M}(h)$ for $0 < h < 1$, together with the two expansions of $\bar{M}(h)$ near the origin and the homoclinic loop L_0, respectively. By (1.3) and (1.4) we have

$$M^+(\frac{h}{2}) = \int_{\widehat{AA_1}} q^+dx - p^+dy, \quad M^-(\frac{h}{2}) = \int_{\widehat{A_1A}} q^-dx - p^-dy = \widetilde{M}^-(\frac{h-1}{2}). \tag{2.1}$$

Then, applying Green's formula twice we obtain

$$\begin{aligned} M^+(\frac{h}{2}) &= -\int\int_{int(\widehat{AA_1}\bigcup\overrightarrow{A_1A})} (p_x^+ + q_y^+)dxdy - \int_{\overrightarrow{AA_1}} p^+(0,y)dy \\ &= -\oint_{\widehat{AA_1}\bigcup\overrightarrow{A_1A}} \overline{p}(x,y)dy - \int_{\overrightarrow{AA_1}} p^+(0,y)dy, \end{aligned}$$

where

$$\overline{p}(x,y) = p^+(x,y) - p^+(0,y) + \int_0^x q_y^+(u,y)du, \tag{2.2}$$

satisfying

$$\overline{p}(0,y) = 0, \quad \overline{p}_x = p_x^+ + q_y^+.$$

Since $\widehat{AA_1}$ can be represented as $x = 1 - \sqrt{h+y^2} \equiv \psi(y,h)$, we have

$$\begin{aligned} M^+(\frac{h}{2}) &= -\Big[\int_{-\sqrt{1-h}}^{\sqrt{1-h}} \overline{p}(\psi(y,h),y)dy + \int_{-\sqrt{1-h}}^{\sqrt{1-h}} p^+(0,y)dy\Big] \\ &\equiv -(I_0(h)+I_1(h)). \end{aligned} \tag{2.3}$$

Note that $p^+(0,y) = \sum\limits_{j=0}^{n} a_{0j}^+ y^j$. It follows that

$$\begin{aligned} I_1(h) &= \int_{-\sqrt{1-h}}^{\sqrt{1-h}} p^+(0,y)dy \\ &= \sum_{j=0}^{n} \frac{a_{0j}^+}{j+1}(1-(-1)^{j+1})(1-h)^{\frac{j+1}{2}} \\ &= \sqrt{1-h}\sum_{k=0}^{[\frac{n}{2}]} \frac{2a_{0,2k}^+}{2k+1}(1-h)^k. \end{aligned} \tag{2.4}$$

By (2.2), we have

$$\begin{aligned} \overline{p}(x,y) &= \sum_{i+j=0}^{n} a_{ij}^+ x^i y^j - \sum_{j=0}^{n} a_{0j}^+ y^j + \sum_{i+j=0}^{n} \frac{jb_{ij}^+}{i+1} x^{i+1} y^{j-1} \\ &\equiv x\sum_{i+j=0}^{n-1} p_{ij}^+ x^i y^j, \end{aligned} \tag{2.5}$$

where

$$p_{ij}^+ = a_{i+1,j}^+ + \frac{j+1}{i+1} b_{i,j+1}^+.$$

In particular, for $n = 2$ we have

$$p_{00}^+ = a_{10}^+ + b_{01}^+,\ p_{10}^+ = a_{20}^+ + \frac{1}{2}b_{11}^+,\ p_{01}^+ = a_{11}^+ + 2b_{02}^+.$$

The definition of $I_0(h)$ and (2.5) yield

$$\begin{aligned} I_0(h) &= \int_{-\sqrt{1-h}}^{\sqrt{1-h}} \overline{p}(\psi(y,h),y)dy \\ &= \sum_{i+j=0}^{n-1} p_{ij}^+ \int_{-\sqrt{1-h}}^{\sqrt{1-h}} (1-\sqrt{h+y^2})^{i+1} y^j dy \\ &= 2\sum_{i+2k=0}^{n-1} p_{i,2k}^+ \int_{0}^{\sqrt{1-h}} (1-\sqrt{h+y^2})^{i+1} y^{2k} dy. \end{aligned} \tag{2.6}$$

For $0 \leq k \leq [\frac{n-1}{2}]$ and $0 \leq r \leq n$ let

$$I_{rk}(h) = \int_0^{\sqrt{1-h}} (h+y^2)^{\frac{r}{2}} y^{2k} dy. \tag{2.7}$$

Then for $0 \leq k \leq [\frac{n-1}{2}]$ and $0 \leq i+2k \leq n-1$

$$\int_0^{\sqrt{1-h}} (1-\sqrt{h+y^2})^{i+1} y^{2k} dy = \sum_{r=0}^{i+1} C_{i+1}^r (-1)^r I_{rk}(h).$$

If $r = 2l,\ l \geq 0$, we see

$$I_{2l,k}(h) = \int_0^{\sqrt{1-h}} (h+y^2)^l y^{2k} dy = \sqrt{1-h}\ \varphi_{l+k}(h), \tag{2.8}$$

where $\varphi_{l+k}(h)$ is a polynomial of degree $l+k$. Hence $I_{2l,k}(h)$ is C^ω at $h=0$. If $r = 2l+1,\ l \geq 0$, then

$$I_{2l+1,k}(h) = \int_0^{\sqrt{1-h}} (h+y^2)^{l+\frac{1}{2}} y^{2k} dy. \tag{2.9}$$

Using the formula

$$\int y^{2k}(h+y^2)^{l+\frac{1}{2}} dy = \frac{y^{2k+1}(h+y^2)^{l+\frac{1}{2}}}{2(l+k+1)} + \frac{(2l+1)h}{2(l+k+1)} \int y^{2k}(h+y^2)^{l-\frac{1}{2}} dy,$$

we have from (2.9) that

$$I_{2l+1,k}(h) = \frac{(1-h)^{k+\frac{1}{2}}}{2(l+k+1)} + \frac{(2l+1)h}{2(l+k+1)} I_{2l-1,k}(h),\ l \geq 1, k \geq 0.$$

It follows that

$$I_{2l+1,k}(h) = (1-h)^{k+\frac{1}{2}} \widetilde{\varphi}_{lk}(h) + \alpha_{lk} h^l I_{1k}(h),\ l \geq 1, k \geq 0, \tag{2.10}$$

where

$$\alpha_{lk} = \frac{(2l+1)(2l-1)\cdots 3}{2^l(l+k+1)(l+k)\cdots(k+2)},$$

and

$$\widetilde{\varphi}_{lk}(h) = \frac{1}{2(l+k+1)} + \frac{(2l+1)h}{2(l+k+1)(l+k)} + \cdots + \frac{(2l+1)(2l-1)\cdots 5 \cdot h^{l-1}}{2^l(l+k+1)(l+k)\cdots(k+2)},$$

which is a polynomial of degree $l-1$ in h. For $k \geq 0$, let

$$\varphi_{lk}^*(h) = \begin{cases} \widetilde{\varphi}_{lk}(h), & l \geq 1, \\ 0, & l = 0, \end{cases} \qquad \alpha_{lk}^* = \begin{cases} \alpha_{lk}, & l \geq 1, \\ 1, & l = 0. \end{cases}$$

By (2.10),

$$I_{2l+1,k}(h) = (1-h)^{k+\frac{1}{2}} \varphi_{lk}^*(h) + \alpha_{lk}^* h^l I_{1k}(h),\ l \geq 0, k \geq 0. \tag{2.11}$$

Further, by using the formula

$$\int y^{2k}(h+y^2)^{\frac{1}{2}}dy = \frac{y^{2k-1}(h+y^2)^{\frac{3}{2}}}{2(k+1)} - \frac{(2k-1)h}{2(k+1)}\int y^{2(k-1)}(h+y^2)^{\frac{1}{2}}dy,$$

we have

$$I_{1k}(h) = \frac{(1-h)^{k-\frac{1}{2}}}{2(k+1)} - \frac{h(2k-1)}{2(k+1)}I_{1,k-1}(h),\ \ k \geq 1.$$

Thus

$$I_{1k}(h) = \sqrt{1-h}\psi_{k-1}(h) + \beta_k h^k I_{10}(h),\ \ k \geq 1, \tag{2.12}$$

where

$$\beta_k = (-1)^k\frac{(2k-1)!!}{2^k(k+1)!},$$

and

$$\psi_{k-1}(h) = \frac{(1-h)^{k-1}}{2(k+1)} - \frac{(2k-1)h(1-h)^{k-2}}{4(k+1)k} + \cdots + (-1)^{k-1}\frac{(2k-1)(2k-3)\cdots 3\cdot h^{k-1}}{2^k(k+1)k(k-1)\cdots 2},$$

which is a polynomial in h of degree $k-1$. Let

$$\psi^*_{k-1}(h) = \begin{cases} \psi_{k-1}(h),\ \ k \geq 1, \\ 0, \qquad\quad k = 0, \end{cases} \qquad \beta^*_k = \begin{cases} \beta_k,\ \ k \geq 1, \\ 1, \quad k = 0. \end{cases}$$

By (2.12),

$$I_{1k}(h) = \sqrt{1-h}\psi^*_{k-1}(h) + \beta^*_k h^k I_{10}(h),\ \ k \geq 0. \tag{2.13}$$

Moreover, by (2.9)

$$I_{10}(h) = \int_0^{\sqrt{1-h}}(h+y^2)^{\frac{1}{2}}dy = \frac{1}{2}[\sqrt{1-h} + h\ \ln(1+\sqrt{1-h}) - \frac{1}{2}h\ \ln h] \equiv -\frac{1}{4}h\ln h + \gamma(h), \tag{2.14}$$

where $\gamma(h)$ is C^ω at $h=0$. We observe that

$$I_{10}(h) = h\varphi_0\Big(\sqrt{\frac{1-h}{h}}\Big),\ \ \varphi_0(u) = \int_0^u \sqrt{1+x^2}dx. \tag{2.15}$$

The function $\varphi_0(u)$ is analytic on R and odd in u.

From (2.11), (2.13) and (2.14), we can obtain

$$\begin{aligned} I_{2l+1,k}(h) &= (1-h)^{k+\frac{1}{2}}\varphi^*_{lk}(h) + \sqrt{1-h}\alpha^*_{lk}\psi^*_{k-1}(h)h^l + \alpha^*_{lk}\beta^*_k h^{l+k}I_{10}(h) \\ &= \sqrt{1-h}\bar{\varphi}_{l+k-1}(h) + \bar{\alpha}_{lk}h^{l+k}I_{10}(h) \\ &= \widetilde{\gamma}(h) - \frac{1}{4}\bar{\alpha}_{lk}h^{l+k+1}\ln h, \end{aligned} \tag{2.16}$$

where $\bar{\varphi}_{l+k-1}(h)$ is a polynomial of degree $l+k-1$, $\bar{\alpha}_{lk}$ are non-zero constants, $\widetilde{\gamma}(h)$ is C^ω at $h=0$, $0 \leq k \leq [\frac{n-1}{2}]$, $0 \leq 2l+1 \leq n$.

Substituting (2.8) and (2.16) into (2.6), and noting that $k+[\frac{i+1}{2}] \leq [\frac{n}{2}]$ and $k+[\frac{i}{2}] \leq [\frac{n-1}{2}]$ since $i+2k \leq n-1$, we have

$$\begin{aligned} I_0(h) &= 2\sum_{i+2k=0}^{n-1} p^{+}_{i,2k}\Big(\sum_{r=0,\ r\ \text{even}}^{i+1} C^r_{i+1}I_{rk} - \sum_{r=1,\ r\ \text{odd}}^{i+1} C^r_{i+1}I_{rk}\Big) \\ &= 2\sum_{i+2k=0}^{n-1} p^{+}_{i,2k}\Big[\sqrt{1-h}\,\nu_{[\frac{i+1}{2}]+k}(h) + \nu_{[\frac{i}{2}]}(h)h^k I_{10})\Big] \\ &= \sqrt{1-h}\,\nu_{[\frac{n}{2}]}(h) + \nu_{[\frac{n-1}{2}]}(h)I_{10} \\ &\equiv \phi_1(h)\ln h + \phi_2(h), \end{aligned} \tag{2.17}$$

where $\nu_j(h)$ denotes a polynomial of h of degree j, $\phi_2(h) \in C^{\omega}$ at $h=0$, $\phi_1(h)$ is a polynomial in h of degree $[\frac{n+1}{2}]$ and $\phi_1(0)=0$.

Now we consider $M^{-}(\frac{h}{2})$ in (2.1). Set $x=\sqrt{-\widetilde{h}}\,\cos\theta,\ y=\sqrt{-\widetilde{h}}\,\sin\theta$ for $-1<\widetilde{h}<0,\ \frac{\pi}{2}\leq\theta\leq\frac{3\pi}{2}$, and let

$$\bar{I}_{ij} = \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} \cos^i\theta\,\sin^j\theta d\theta = (-1)^{i+j}\int_{\frac{-\pi}{2}}^{\frac{\pi}{2}} \cos^i\theta\,\sin^j\theta d\theta = (-1)^{i+j}\Gamma_{ij},$$

where

$$\Gamma_{ij} = \begin{cases} 0,\ j\ \text{odd}, \\ \text{a positive constant},\ j\ \text{even}. \end{cases}$$

It follows from (1.3) that

$$\begin{aligned} \widetilde{M}^{-}(\frac{\widetilde{h}}{2}) &= \int_{H^{-}(x,y)=\frac{\widetilde{h}}{2}} q^{-}dx - p^{-}dy \\ &= -\sum_{i+j=0}^{n}(-\widetilde{h})^{\frac{i+j+1}{2}}\int_{\frac{\pi}{2}}^{\frac{3\pi}{2}}(b^{-}_{ij}\cos^i\theta\,\sin^{j+1}\theta + a^{-}_{ij}\cos^{i+1}\theta\,\sin^j\theta)d\theta \\ &= -\sum_{i+j=0}^{n}(-\widetilde{h})^{\frac{i+j+1}{2}}(b^{-}_{ij}\bar{I}_{i,j+1} + a^{-}_{ij}\bar{I}_{i+1,j}) \\ &= -\sum_{l=0}^{n}(-\widetilde{h})^{\frac{l+1}{2}}\sum_{j=0}^{l}(b^{-}_{l-j,j}\bar{I}_{l-j,j+1} + a^{-}_{l-j,j}\bar{I}_{l-j+1,j}) \\ &= -\sum_{l=0}^{n} e_l(-\widetilde{h})^{\frac{l+1}{2}}, \end{aligned} \tag{2.18}$$

where

$$\begin{aligned} e_l &= \sum_{j=0}^{l}(b^-_{l-j,j}\bar{I}_{l-j,j+1} + a^-_{l-j,j}\bar{I}_{l-j+1,j}) \\ &= (-1)^{l+1}\sum_{j=0}^{l}(b^-_{l-j,j}\Gamma_{l-j,j+1} + a^-_{l-j,j}\Gamma_{l-j+1,j}) \qquad (2.19) \\ &= (-1)^{l+1}\Big[\sum_{j=0,\ j\ \text{even}}^{l} a^-_{l-j,j}\Gamma_{l-j+1,j} + \sum_{j=1,\ j\ \text{odd}}^{l} b^-_{l-j,j}\Gamma_{l-j,j+1}\Big],\ 0 \le l \le n. \end{aligned}$$

By (2.1) and (2.18),

$$M^-(\frac{h}{2}) = -\sqrt{1-h}\sum_{l=0}^{n} e_l(1-h)^{\frac{l}{2}}, \quad 0 < h < 1, \tag{2.20}$$

where for $n = 2$

$$e_0 = -2a^-_{00},\ e_1 = \frac{\pi}{2}(a^-_{10} + b^-_{01}),\ e_2 = -\frac{2}{3}(2a^-_{20} - b^-_{11} + a^-_{02}).$$

Hence by (1.10), (2.1), (2.3), (2.4), (2.17) and (2.20) we obtain the following lemma.

Lemma 2.1. *For system* (1.9), *the first order Melnikov function has the following form*

$$\bar{M}(h) = \sqrt{1-h}f_n(\sqrt{1-h}) + g_{[\frac{n-1}{2}]}(1-h)I_{10}(h),$$

where f_n and $g_{[\frac{n-1}{2}]}$ are polynomials of degrees n and $[\frac{n-1}{2}]$ respectively.

In the following we study the expansions of $\bar{M}(h)$ near $h = 0, 1$. By (2.14) and Lemma 2.1, the lemma below holds immediately.

Lemma 2.2. *For system* (1.9) *the first order Melnikov function $\bar{M}(h)$ has the following expansion near the homoclinic loop L_0*

$$\bar{M}(h) = (\sum_{i=1}^{[\frac{n+1}{2}]} b^*_i h^i)\ \ln h + \sum_{j\ge 0} b_j h^j, \quad 0 < h \ll 1,$$

where for $n = 2$

$$b^*_1 = -\tfrac{1}{2}(\ a^+_{10} +\ b^+_{01} + 2a^+_{20} +\ b^+_{11}),$$

$$b_0 = -a^+_{10} - b^+_{01} - \tfrac{2}{3}\ a^+_{20} - \tfrac{1}{3}\ b^+_{11} - 2\ a^+_{00} - \tfrac{2}{3}\ a^+_{02} + 2a^-_{00} - \tfrac{\pi}{2}a^-_{10} - \tfrac{\pi}{2}b^-_{01} + \tfrac{4}{3}a^-_{20} - \tfrac{3}{2}b^-_{11} + \tfrac{2}{3}a^-_{02},$$

$$\begin{aligned} b_1 = {} & (\tfrac{1}{2} + \ln 2)\ a^+_{10} + (\tfrac{1}{2} + \ln 2)b^+_{01} + (2\ln 2 - 1)a^+_{20} + (\ln 2 - \tfrac{1}{2})\ b^+_{11} + a^+_{00} + a^+_{02} - a^-_{00} \\ & + \tfrac{\pi}{2}a^-_{10} + \tfrac{\pi}{2}b^-_{01} - 2a^-_{20} + b^-_{11} - a^-_{02}, \end{aligned}$$

$$b_2 = \tfrac{1}{4}\ a^+_{20} + \tfrac{1}{8}\ b^+_{11} - \tfrac{1}{4}\ a^+_{02} - \tfrac{1}{8}\ a^+_{10} - \tfrac{1}{8}\ b^+_{01} + \tfrac{1}{4}\ a^+_{00} - \tfrac{1}{4}a^-_{00} + \tfrac{1}{2}a^-_{02} - \tfrac{1}{4}b^-_{11} + \tfrac{1}{4}a^-_{02},$$

……

Noticing that in (2.15) $\varphi_0(u) \in C^w$ on R and is odd in u, we can write for $|u|$ small

$$\varphi_0(u) = \sum_{i=0}^{\infty} \varsigma_i u^{2i+1},$$

where ς_i is a constant, $i \geq 0$. Then, it follows from (2.15) for $1-h>0$ small

$$I_{10}(h) = h\varphi_0\Big(\sqrt{\frac{1-h}{h}}\Big) = \sqrt{h(1-h)}\sum_{i=0}^{\infty} \varsigma_i (\frac{1-h}{h})^i \equiv \sqrt{1-h}\phi_3(1-h), \tag{2.21}$$

where $\phi_3(u) \in C^w$ at $u=0$. Following (2.21) and Lemma 2.1, we have

Lemma 2.3. *For system (1.9) the first order Melnikov function $\bar{M}(h)$ has the following expansion near the origin*

$$\bar{M}(h) = \sqrt{1-h}\big[\sum_{i=0}^{n} c_i (1-h)^{\frac{i}{2}} + \sum_{j\geq 1} c_{n+j}(1-h)^{[\frac{n}{2}]+j}\big], \quad 0 < 1-h \ll 1,$$

where for $n=2$

$$c_0 = 2(a_{00}^- - a_{00}^+), \ c_1 = -\tfrac{\pi}{2}(a_{10}^- + b_{01}^-),$$

$$c_2 = -\tfrac{2}{3}(a_{02}^+ + b_{01}^+ + a_{10}^+ - 2a_{20}^- + b_{11}^- - a_{02}^-),$$

$$c_3 = -\tfrac{2}{15}(2a_{20}^+ + a_{10}^+ + b_{11}^+ + b_{01}^+), \cdots\cdots$$

3 Proof of the main results

Proof of Theorem 1.1. (1) For simplicity, we let $p^{\pm}$ and $q^{\pm}$ in (1.8) satisfy

$$p^{\pm}(x,y) = \sum_{i=0}^{n} a_i^{\pm} x^i, \ q^{\pm}(x,y) \equiv 0, \tag{3.1}$$

that is, $p^{\pm}(x,y)$ are univariate polynomials of the variable x. By (2.4) and (2.5), we have

$$I_1(h) = 2a_0^+\sqrt{1-h}, \ \ \overline{p}(x,y) = \sum_{i=1}^{n} a_i^+ x^i,$$

From (1.6), the equation $H^+(x,y) = \frac{h}{2}$ is equivalent to

$$2x - x^2 = \omega, \tag{3.2}$$

where $\omega = 1-h-y^2$. Consider equation (3.2) in x near $x=0$. There is a unique C^∞ solution

$$x = \sigma(\omega) = \frac{1}{2}\omega + O(\omega^2) \equiv \sum_{j\geq 1} g_j \omega^j, \tag{3.3}$$

where

$$g_1 = \frac{1}{2},\ g_2 = \frac{1}{8},\ g_3 = \frac{1}{16},\ g_4 = \frac{5}{128}, \cdots$$

Then, (3.3) implies that the arc $\widehat{AA_1}$ can be represented by $x = \sigma(\omega)$ near the origin. That is, in this case the function $\psi(y, h)$ in (2.3) can be taken as $x = \sigma(\omega)$ with $\omega = 1 - h - y^2,\ 1 - h > 0$ small. Note that

$$\overline{p}(\sigma(\omega), y) = \sum_{i=1}^{n} a_i^+ (\sigma(\omega))^i = \sum_{j\geq 1} \widetilde{p}_j \omega^j,$$

where $\widetilde{p}_1 = \frac{1}{2}a_1^+$, $\widetilde{p}_2 = \frac{1}{4}a_2^+ + \frac{1}{8}a_1^+,\ \cdots,\ \widetilde{p}_n = \frac{a_n^+}{2^n} + L(a_1^+, \cdots, a_{n-1}^+), \cdots$, throughout this paper $L(\cdot)$ denotes a linear combination. Then by (2.6), we have for $0 < 1 - h \ll 1$

$$\begin{aligned} I_0(h) &= \sum_{j\geq 1} \int_{-\sqrt{1-h}}^{\sqrt{1-h}} \widetilde{p}_j \omega^j dy \\ &= 2\sum_{j\geq 1} \widetilde{p}_j \int_0^{\sqrt{1-h}} (1 - h - y^2)^j dy \\ &= \sqrt{1-h} \sum_{j\geq 1} \widetilde{p}_j B_j (1-h)^j, \end{aligned}$$

where $B_j = 2\int_0^1 (1-u^2)^j du$ is a positive constant for $j \geq 1$. Hence, by (2.3)

$$\begin{aligned} M^+(\frac{h}{2}) &= -\big(I_0(h) + I_1(h)\big) \\ &= -\big(\sqrt{1-h}\sum_{j\geq 1} \widetilde{p}_j B_j (1-h)^j + 2a_0^+\sqrt{1-h}\big) \\ &\equiv \sqrt{1-h}\sum_{j\geq 0} c_j^* (1-h)^j,\ \text{for } 0 < 1 - h \ll 1, \end{aligned} \tag{3.4}$$

where $c_j^* = \begin{cases} -2a_0^+,\ j = 0, \\ -\widetilde{p}_j B_j,\ \ j \geq 1. \end{cases}$ Under (3.1), we also have from (2.19) and (2.20)

$$M^-(\frac{h}{2}) = -\sqrt{1-h}\sum_{l=0}^{n} e_l (1-h)^{\frac{l}{2}},\quad 0 < h < 1, \tag{3.5}$$

where $e_l = (-1)^{l+1} a_l^- \Gamma_{l+1,0}$, $\Gamma_{l+1,0}$ is a positive constant, $0 \leq l \leq n$. Thus, by (1.10), (3.4) and (3.5)

$$\bar{M}(h) = \sqrt{1-h}\Big[\sum_{i=0}^{n} c_i (1-h)^{\frac{i}{2}} + \sum_{j\geq 1} c_{n+j} (1-h)^{[\frac{n}{2}]+j}\Big],\quad 0 < 1 - h \ll 1, \tag{3.6}$$

with

$$c_0 = c_0^* - e_0 = -2a_0^+ + \Gamma_{10} a_0^-, \qquad c_1 = -e_1 = -\Gamma_{20} a_1^-,$$

$$c_2 = c_1^* - e_2 = -\frac{B_1}{2} a_1^+ + \Gamma_{30} a_2^-, \qquad c_3 = -e_3 = -\Gamma_{40} a_3^-, \quad \cdots\cdots$$

$$c_n = \begin{cases} c^*_{\frac{n}{2}} - e_n = -\frac{B_{\frac{n}{2}}}{2^{\frac{n}{2}}} a^+_{\frac{n}{2}} + (-1)^n \Gamma_{n+1,0} a_n^- + L(a_1^+, \cdots, a^+_{\frac{n}{2}-1}),\ n \text{ even}, \\ \\ -e_n = (-1)^n \Gamma_{n+1,0} a_n^-,\ n \text{ odd}, \end{cases}$$

$$c_{n+1} = c^*_{[\frac{n}{2}]+1} = -\frac{B_{[\frac{n}{2}]+1}}{2^{[\frac{n}{2}]+1}} a^+_{[\frac{n}{2}]+1} + L(a_1^+, \cdots, a^+_{[\frac{n}{2}]}),$$

$$\cdots\cdots$$

$$c_{2n-[\frac{n}{2}]} = c_n^* = -\frac{B_n}{2^n} a_n^+ + L(a_1^+, \cdots, a_{n-1}^+),$$

$$\cdots\cdots$$

Following the formulas above, we have

$$\frac{\partial(c_0, c_1, c_2, \cdots, c_n, c_{n+1}, \cdots, c_{2n-[\frac{n}{2}]})}{\partial(a_0^+, a_1^+, \cdots, a^+_{[\frac{n}{2}]}, a^+_{[\frac{n}{2}]+1}, \cdots, a_n^+, a_0^-, a_1^-, \cdots, a_n^-)} =$$

$$\begin{pmatrix} -2 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 & \Gamma_{10} & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 & 0 & -\Gamma_{20} & \cdots & 0 \\ 0 & -\frac{B_1}{2} & \cdots & 0 & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & * & \cdots & X^* & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & (-1)^n\Gamma_{n+1,0} \\ 0 & * & \cdots & * & -\frac{B_{[\frac{n}{2}]+1}}{2^{[\frac{n}{2}]+1}} & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ 0 & * & \cdots & * & * & -\frac{B_{[\frac{n}{2}]+2}}{2^{[\frac{n}{2}]+2}} & \cdots & 0 & 0 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & * & \cdots & * & * & * & \cdots & -\frac{B_n}{2^n} & 0 & 0 & \cdots & 0 \end{pmatrix}$$

where

$$X^* = \begin{cases} -\frac{B_{\frac{n}{2}}}{2^{\frac{n}{2}}},\ n \text{ even}, \\ 0,\ n \text{ odd}. \end{cases}$$

Clearly, the rank of this matrix is $2n - [\frac{n}{2}] + 1$, which means that $c_i,\ 0 \le i \le 2n - [\frac{n}{2}]$ can be chosen as free parameters such that $0 < |c_0| \ll |c_1| \ll |c_2| \ll \cdots \ll |c_n| \ll |c_{n+1}| \ll |c_{n+2}| \ll \cdots \ll |c_{2n-[\frac{n}{2}]}| \ll 1$, and $c_i c_{i+1} < 0,\ 0 \le i \le 2n - [\frac{n}{2}] - 1$. Hence, by (3.6), in this case $\bar{M}(h)$ has $2n - [\frac{n}{2}]$ positive simple zeros in $(0, 1)$ near $h = 1$. Therefore, system (1.9) can have $2n - [\frac{n}{2}]$ $(=n + [\frac{n+1}{2}],\ n \ge 1)$ limit cycles near the origin, which means $N_{Hopf}(n) \ge n + [\frac{n+1}{2}]$.

(2) Here, we suppose $p^{\pm}$ and $q^{\pm}$ in (1.8) satisfy

$$p^+(x,y) = \sum_{i=0}^{n} a_i^+ y^i,\ \ q^+(x,y) = \sum_{i=0}^{n} b_i^+ y^i,\ \ p^-(x,y) = \sum_{i=0}^{n} a_i^- x^i,\ \ q^-(x,y) \equiv 0, \tag{3.7}$$

where $p^+(x,y)$ and $q^+(x,y)$ are independent of x and $p^-(x,y)$ is independent of y . Then, from (2.4) to (2.14), we have

$$\overline{p}(x,y) = x\sum_{i=1}^{n} b_i^+ i y^{i-1} = x\sum_{j=0}^{n-1} p_{0j}^+ y^j,$$

where

$$p_{00}^+ = b_1^+, p_{01}^+ = 2b_2^+, \cdots, p_{0,n-1}^+ = nb_n^+. \tag{3.8}$$

Moreover,

$$I_1(h) = \sqrt{1-h}\sum_{k=0}^{[\frac{n}{2}]} \frac{2a_{2k}^+}{2k+1}(1-h)^k,$$

and

$$\begin{aligned}
I_0(h) &= \sum_{j=0}^{n-1} p_{0j}^+ \int_{-\sqrt{1-h}}^{\sqrt{1-h}} (1-\sqrt{h+y^2})y^j dy \\
&= 2\sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^+ \int_0^{\sqrt{1-h}} (1-\sqrt{h+y^2})y^{2k} dy \\
&= 2\Big[\sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^+ \int_0^{\sqrt{1-h}} y^{2k} dy - \sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^+ \int_0^{\sqrt{1-h}} \sqrt{h+y^2} y^{2k} dy\Big] \\
&= 2\Big[\sqrt{1-h}\sum_{k=0}^{[\frac{n-1}{2}]} \frac{p_{0,2k}^+}{2k+1}(1-h)^k - \sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^+ I_{1k}(h)\Big],
\end{aligned}$$

where I_{1k} is given by (2.7). Hence by (2.3), (2.13), (2.14) and the above

$$\begin{aligned}
\frac{M^+(\frac{h}{2})}{-2\sqrt{1-h}} &= \sum_{k=0}^{[\frac{n}{2}]} \frac{a_{2k}^+}{2k+1}(1-h)^k + \sum_{k=0}^{[\frac{n-1}{2}]} \frac{p_{0,2k}^+}{2k+1}(1-h)^k - \sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^+ \psi_{k-1}^*(h) \\
&\quad - (1-h)^{-\frac{1}{2}} \sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^+ \beta_k^* h^k I_{10}(h)
\end{aligned}$$

$$
\begin{aligned}
&= \sum_{k=0}^{[\frac{n}{2}]} \frac{a_{2k}^{+}}{2k+1}(1-h)^k + \sum_{k=0}^{[\frac{n-1}{2}]} \frac{p_{0,2k}^{+}}{2k+1}(1-h)^k - \sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^{+}\psi_{k-1}^{*}(h) \\
&\quad - \frac{1}{2}\sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^{+}\beta_k^{*}h^k - \frac{1}{2}\sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^{+}\beta_k^{*}h^{k+1}(1-h)^{-\frac{1}{2}}\ln(1+\sqrt{1-h}) \\
&\quad + \frac{1}{4}\sum_{k=0}^{[\frac{n-1}{2}]} p_{0,2k}^{+}\beta_k^{*}(1-h)^{-\frac{1}{2}}h^{k+1}\ln h \\
&\equiv \sum_{k=0}^{\infty} v_k h^k + \sum_{k=0}^{\infty} v_k^{*} h^{k+1}\ln h, \quad \text{for } 0 < h \ll 1,
\end{aligned}
\tag{3.9}
$$

where

$$
\begin{aligned}
& v_0 = a_0^{+} + L(a_2^{+}, \cdots, a_{2[\frac{n}{2}]}^{+}, p_{00}^{+}, \cdots, p_{0,2[\frac{n-1}{2}]}^{+}), \\
& v_1 = -\frac{a_2^{+}}{3} + L(a_4^{+}, \cdots, a_{2[\frac{n}{2}]}^{+}, p_{00}^{+}, \cdots, p_{0,2[\frac{n-1}{2}]}^{+}), \\
& v_2 = \frac{a_4^{+}}{5} + L(a_6^{+}, \cdots, a_{2[\frac{n}{2}]}^{+}, p_{00}^{+}, \cdots, p_{0,2[\frac{n-1}{2}]}^{+}), \quad \cdots\cdots \\
& v_{[\frac{n}{2}]} = \frac{a_{2[\frac{n}{2}]}^{+}}{2[\frac{n}{2}]+1}(-1)^{[\frac{n}{2}]} + L(p_{00}^{+}, \cdots, p_{0,2[\frac{n-1}{2}]}^{+}), \\
& v_0^{*} = \frac{1}{4}p_{00}^{+}, \ \ v_1^{*} = \frac{1}{4}p_{02}^{+}\beta_1^{*} + L(p_{00}^{+}), \\
& v_2^{*} = \frac{1}{4}p_{04}^{+}\beta_2^{*} + L(p_{00}^{+}, p_{02}^{+}), \\
& \cdots\cdots \\
& v_{[\frac{n-1}{2}]}^{*} = \frac{1}{4}p_{0,2[\frac{n-1}{2}]}^{+}\beta_{[\frac{n-1}{2}]}^{*} + L(p_{00}^{+}, p_{02}^{+}, \cdots, p_{0,2[\frac{n-1}{2}]-2}^{+}).
\end{aligned}
\tag{3.10}
$$

Moreover, from (3.5) we have for $0 < h \ll 1$

$$
\frac{M^{-}(\frac{h}{2})}{-2\sqrt{1-h}} = \frac{1}{2}\sum_{l=0}^{n} e_l(1-h)^{\frac{l}{2}} = \frac{1}{2}\big[\sum_{i=0}^{[\frac{n}{2}]} e_{2i}(1-h)^i + \sqrt{1-h}\sum_{j=0}^{[\frac{n-1}{2}]} e_{2j+1}(1-h)^j\big] \equiv C_1 + C_2,
\tag{3.11}
$$

where

$$\begin{aligned} C_1 &= \tfrac{1}{2}\sum_{i=0}^{[\frac{n}{2}]} e_{2i}(1-h)^i = -\tfrac{1}{2}\sum_{i=0}^{[\frac{n}{2}]} a_{2i}^{-}\Gamma_{2i+1,0}(1-h)^i \\ &= L(a_0^-, a_2^-, \cdots, a_{2[\frac{n}{2}]}^-) + L(a_2^-, a_4^-, \cdots, a_{2[\frac{n}{2}]}^-)h + \cdots - \tfrac{1}{2}a_{2[\frac{n}{2}]}^-\Gamma_{2[\frac{n}{2}]+1,0}(-1)^{[\frac{n}{2}]}h^{[\frac{n}{2}]}, \end{aligned} \tag{3.12}$$

$$\begin{aligned} 2C_2 &= \sqrt{1-h}\sum_{j=0}^{[\frac{n-1}{2}]} e_{2j+1}(1-h)^j = \sqrt{1-h}\sum_{j=0}^{[\frac{n-1}{2}]} a_{2j+1}^-\Gamma_{2j+2,0}(1-h)^j \\ &= \sqrt{1-h}\sum_{j=0}^{[\frac{n-1}{2}]} a_{2j+1}^-\Gamma_{2j+2,0}\sum_{k=0}^{j} C_j^k(-1)^k h^k \\ &= \sqrt{1-h}\sum_{k=0}^{[\frac{n-1}{2}]}\Big(\sum_{j=k}^{[\frac{n-1}{2}]} a_{2j+1}^-\Gamma_{2j+2,0}C_j^k(-1)^k\Big)h^k. \end{aligned} \tag{3.13}$$

Let

$$\sqrt{1-h} = \sum_{i\geq 0}\mu_i h^i,\ 0<h\ll 1, \tag{3.14}$$

where

$$\mu_0 = 1,\ \mu_1 = -\frac{1}{2},\ \mu_i = -\frac{(2i-3)!!}{2^i},\ i\geq 2. \tag{3.15}$$

Denote

$$D_k = \sum_{m=k}^{[\frac{n-1}{2}]} a_{2m+1}^-\Gamma_{2m+2,0}C_m^k(-1)^k,\ 0\leq k\leq [\frac{n-1}{2}]. \tag{3.16}$$

Then, by (3.13), (3.14) and (3.16) we have

$$2C_2 = \sum_{i=0}^{[\frac{n-1}{2}]}(\sum_{j=0}^{i}\mu_{i-j}D_j)h^i + \sum_{k\geq 1}\Big(\sum_{s=0}^{[\frac{n-1}{2}]}\mu_{k+s}D_{[\frac{n-1}{2}]-s}\Big)h^{[\frac{n-1}{2}]+k},\ 0<h\ll 1. \tag{3.17}$$

From (3.11), (3.12), (3.16) and (3.17), we get

$$\frac{M^-(\frac{h}{2})}{-2\sqrt{1-h}} = \sum_{i\geq 0}\omega_i h^i,\ 0<h\ll 1, \tag{3.18}$$

where

$$\omega_i = L(a_0^-, a_1^-, \cdots, a_n^-),\ 0 \le i \le [\tfrac{n-1}{2}],$$

$$\omega_{[\frac{n-1}{2}]+1} = \begin{cases} \frac{1}{2}\sum\limits_{s=0}^{\frac{n-1}{2}} \mu_{1+s} D_{\frac{n-1}{2}-s},\ n \text{ odd}, \\ \\ \frac{1}{2}\sum\limits_{s=0}^{\frac{n-2}{2}} \mu_{1+s} D_{\frac{n-2}{2}-s} + \frac{(-1)^{\frac{n}{2}+1}}{2}\Gamma_{n+1,0}a_n^-,\ n \text{ even}, \end{cases} \tag{3.19}$$

$$\omega_{[\frac{n-1}{2}]+k} = \frac{1}{2}\sum_{s=0}^{[\frac{n-1}{2}]} \mu_{k+s} D_{[\frac{n-1}{2}]-s},\ k \ge 2.$$

Hence, it follows from (3.9) and (3.18) that

$$\frac{M(\frac{h}{2})}{-2\sqrt{1-h}} = \sum_{i\ge 0}(\widetilde{v}_i h^i + v_i^* h^{i+1}\ln h),\ 0 < h \ll 1, \tag{3.20}$$

where by (3.10) and (3.19)

$$
\begin{aligned}
&\widetilde{v}_0 = a_0^+ + L(a_2^+, \cdots, a_{2[\frac{n}{2}]}^+, p_{00}^+, \cdots, p_{0,2[\frac{n-1}{2}]}^+) + L(a_0^-, a_1^-, \cdots, a_n^-),\\
&v_0^* = \tfrac{1}{4} p_{00}^+,\\
&\widetilde{v}_1 = -\tfrac{a_2^+}{3} + L(a_4^+, \cdots, a_{2[\frac{n}{2}]}^+, p_{00}^+, \cdots, p_{0,2[\frac{n-1}{2}]}^+) + L(a_0^-, a_1^-, \cdots, a_n^-),\\
&v_1^* = \tfrac{1}{4} p_{02}^+ \beta_1^* + L(p_{00}^+),\\
&\cdots\cdots\\
&\widetilde{v}_{[\frac{n-1}{2}]} = \begin{cases} \frac{a_{n-1}^+}{n}(-1)^{\frac{n-1}{2}} + L(p_{00}^+, p_{02}^+, \cdots, p_{0,n-1}^+) + L(a_0^-, a_1^-, \cdots, a_n^-), \ n \text{ odd},\\[2ex] \frac{a_{n-2}^+}{n-1}(-1)^{\frac{n}{2}-1} + L(a_n^+, p_{00}^+, p_{02}^+, \cdots, p_{0,n-2}^+) + L(a_0^-, a_1^-, \cdots, a_n^-), \ n \text{ even}, \end{cases}\\[2ex]
&v_{[\frac{n-1}{2}]}^* = \tfrac{1}{4} p_{0,2[\frac{n-1}{2}]}^+ \beta_{[\frac{n-1}{2}]}^* + L(p_{00}^+, p_{02}^+, \cdots, p_{0,2[\frac{n-1}{2}]-2}^+),\\[2ex]
&\widetilde{v}_{[\frac{n-1}{2}]+1} = \begin{cases} \frac{1}{2} \sum\limits_{s=0}^{\frac{n-1}{2}} \mu_{1+s} D_{\frac{n-1}{2}-s} + L(p_{00}^+, p_{02}^+, \cdots, p_{0,n-1}^+), \ n \text{ odd},\\[2ex] \frac{a_n^+}{n+1}(-1)^{\frac{n}{2}} + L(p_{00}^+, p_{02}^+, \cdots, p_{0,n-2}^+) + L(a_n^-, a_1^-, a_3^-, \cdots, a_{n-1}^-), \ n \text{ even}, \end{cases}\\
&v_{[\frac{n-1}{2}]+i}^* = O(|v_0^*, v_1^*, \cdots, v_{[\frac{n-1}{2}]}^*|), \ i \geq 1, (\text{obtained by } (3.9) \ \text{ and } (3.10)),\\
&\widetilde{v}_{[\frac{n-1}{2}]+j} = \tfrac{1}{2} \sum_{s=0}^{[\frac{n-1}{2}]} \mu_{j+s} D_{[\frac{n-1}{2}]-s} + L(p_{00}^+, p_{02}^+, \cdots, p_{0,2[\frac{n-1}{2}]}^+), \ j \geq 2.
\end{aligned}
\tag{3.21}
$$

If n is odd, by the formulas of $\widetilde{v}_{[\frac{n-1}{2}]+j}, \ j \geq 1$ in the above and $D_k, \ 0 \leq k \leq [\frac{n-1}{2}]$ in (3.16) we have

$$
\begin{aligned}
\widetilde{v}_{\frac{n-1}{2}+j} &= \tfrac{1}{2} \sum_{s=0}^{\frac{n-1}{2}} \mu_{j+s} D_{\frac{n-1}{2}-s} + L(p_{00}^+, p_{02}^+, \cdots, p_{0,n-1}^+)\\
&= \tfrac{1}{2} \sum_{s=0}^{\frac{n-1}{2}} \sum_{m=\frac{n-1}{2}-s}^{\frac{n-1}{2}} \mu_{j+s} a_{2m+1}^- \Gamma_{2m+2,0} C_m^{\frac{n-1}{2}-s} (-1)^{\frac{n-1}{2}-s} + L(p_{00}^+, p_{02}^+, \cdots, p_{0,n-1}^+)
\end{aligned}
$$

$$= \frac{1}{2}\sum_{m=0}^{\frac{n-1}{2}}\Big(\sum_{s=\frac{n-1}{2}-m}^{\frac{n-1}{2}}\mu_{j+s}\Gamma_{2m+2,0}C_m^{\frac{n-1}{2}-s}(-1)^{\frac{n-1}{2}-s}\Big)a^-_{2m+1} + L(p^+_{00}, p^+_{02}, \cdots, p^+_{0,n-1})$$

$$= \frac{1}{2}\sum_{m=0}^{\frac{n-1}{2}}\Big(\sum_{t=0}^{m}\mu_{j+\frac{n-1}{2}-t}\Gamma_{2m+2,0}C_m^{t}(-1)^{t}\Big)a^-_{2m+1} + L(p^+_{00}, p^+_{02}, \cdots, p^+_{0,n-1}), \tag{3.22}$$

where $j \geq 1$. For $1 \leq j \leq \frac{n+1}{2}$, it follows from (3.22)

$$A_1 \equiv \frac{\partial(\widetilde{v}_{\frac{n-1}{2}+1}, \widetilde{v}_{\frac{n-1}{2}+2}, \cdots, \widetilde{v}_n)}{\partial(a^-_1, a^-_3, \cdots, a^-_{n-2}, a^-_n)} \tag{3.23}$$

$$= \begin{pmatrix} \frac{1}{2}\mu_{1+\frac{n-1}{2}}\Gamma_{20} & \frac{1}{2}\sum\limits_{t=0}^{1}\mu_{1+\frac{n-1}{2}-t}\Gamma_{40}C_1^t(-1)^t & \cdots & \frac{1}{2}\sum\limits_{t=0}^{\frac{n-1}{2}}\mu_{1+\frac{n-1}{2}-t}\Gamma_{n+1,0}C^t_{\frac{n-1}{2}}(-1)^t \\ \frac{1}{2}\mu_{2+\frac{n-1}{2}}\Gamma_{20} & \frac{1}{2}\sum\limits_{t=0}^{1}\mu_{2+\frac{n-1}{2}-t}\Gamma_{40}C_1^t(-1)^t & \cdots & \frac{1}{2}\sum\limits_{t=0}^{\frac{n-1}{2}}\mu_{2+\frac{n-1}{2}-t}\Gamma_{n+1,0}C^t_{\frac{n-1}{2}}(-1)^t \\ \cdots & \cdots & \cdots & \cdots \\ \frac{1}{2}\mu_n\Gamma_{20} & \frac{1}{2}\sum\limits_{t=0}^{1}\mu_{n-t}\Gamma_{40}C_1^t(-1)^t & \cdots & \frac{1}{2}\sum\limits_{t=0}^{\frac{n-1}{2}}\mu_{n-t}\Gamma_{n+1,0}C^t_{\frac{n-1}{2}}(-1)^t \end{pmatrix}.$$

Let c_i denote the ith column of A_1, $a \times c_i$ (or c_i/a) indicate multiplying (or dividing) each element of the ith column by a, and $c_i + b \times c_j$ indicate that each element of the jth column times b adds to the according element of the ith column, a and b are constants. We make elementary transformations to A_1 with the following steps

S_1. $c_i/(\frac{1}{2}\Gamma_{2i,0})$, $i = 1, 2, \cdots, \frac{n+1}{2}$.

S_2. $c_i + (-1) \times c_1$, $i = 2, 3, \cdots, \frac{n+1}{2}$.

S_3. $c_i + (-C^1_{i-1}) \times c_2$, $i = 3, 4, \cdots, \frac{n+1}{2}$; $(-1) \times c_2$.

……

$S_{\frac{n+1}{2}}$. $c_{\frac{n+1}{2}} + (-\frac{n-1}{2}) \times c_{\frac{n-1}{2}}$, $c_{\frac{n-1}{2}}/(-1)^{\frac{n-3}{2}}$ and $c_{\frac{n+1}{2}}/(-1)^{\frac{n-1}{2}}$.

Then A_1 becomes

$$\tilde{A}_1 \equiv \begin{pmatrix} \mu_{\frac{n-1}{2}+1} & \mu_{\frac{n-1}{2}} & \cdots & \mu_2 & \mu_1 \\ \mu_{\frac{n-1}{2}+2} & \mu_{\frac{n-1}{2}+1} & \cdots & \mu_3 & \mu_2 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ \mu_n & \mu_{n-1} & \cdots & \mu_{\frac{n-1}{2}+2} & \mu_{\frac{n-1}{2}+1} \end{pmatrix}. \tag{3.24}$$

By (3.15), we can prove $\mathrm{Rank}(\tilde{A}_1) = \frac{n+1}{2}$. See Appendix for the proof. Therefore, by using (3.21) and (3.24) we obtain

$$\frac{\partial(\widetilde{v}_0, \widetilde{v}_1, \cdots, \widetilde{v}_{\frac{n-1}{2}}, v^*_0, v^*_1, \cdots, v^*_{\frac{n-1}{2}}, \widetilde{v}_{\frac{n-1}{2}+1}, \cdots, \widetilde{v}_n)}{\partial(a^+_0, a^+_2, \cdots, a^+_{n-1}, p^+_{00}, p^+_{02}, \cdots, p^+_{0,n-1}, a^-_1, a^-_3, \cdots, a^-_n, a^-_0, a^-_2, \cdots, a^-_{n-1})} =$$

$$\begin{pmatrix}
1 & \ast & \cdots & \ast & \ast & \ast & \cdots & \ast & \ast & \ast & \cdots & \ast & \ast & \ast & \cdots & \ast \\
0 & -\frac{1}{3} & \cdots & \ast & \ast & \ast & \cdots & \ast & \ast & \ast & \cdots & \ast & \ast & \ast & \cdots & \ast \\
\cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\
0 & 0 & \cdots & \frac{(-1)^{\frac{n-1}{2}}}{n} & \ast & \ast & \cdots & \ast & \ast & \ast & \cdots & \ast & \ast & \ast & \cdots & \ast \\
0 & 0 & \cdots & 0 & \frac{1}{4} & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\
0 & 0 & \cdots & 0 & \ast & \frac{\beta_1^*}{4} & \cdots & 0 & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\
\cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\
0 & 0 & \cdots & 0 & \ast & \ast & \cdots & \frac{\beta^*_{\frac{n-1}{2}}}{4} & 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\
0 & 0 & \cdots & 0 & \ast & \ast & \cdots & \ast & & & & & 0 & 0 & \cdots & 0 \\
\cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & & A_1 & & & \cdots & \cdots & \cdots & \cdots \\
0 & 0 & \cdots & 0 & \ast & \ast & \cdots & \ast & & & & & 0 & 0 & \cdots & 0
\end{pmatrix}$$

It is easy to see that the rank of this matrix is $n + \frac{n-1}{2} + 2$. Thus, we can choose $\widetilde{v}_0, v_0^*, \widetilde{v}_1, v_1^*, \cdots, \widetilde{v}_{\frac{n-1}{2}}, v^*_{\frac{n-1}{2}}, \widetilde{v}_{\frac{n-1}{2}+1} \cdots, \widetilde{v}_n$ as free parameters such that

$$0 < \widetilde{v}_0 \ll v_0^* \ll \widetilde{v}_1 \ll v_1^* \ll \cdots \ll \widetilde{v}_{\frac{n-1}{2}} \ll v^*_{\frac{n-1}{2}}$$

$$\ll \widetilde{v}_{\frac{n-1}{2}+1} \ll -\widetilde{v}_{\frac{n-1}{2}+2} \ll \cdots \ll (-1)^{\frac{n-1}{2}} \widetilde{v}_n \ll 1,$$

or

$$0 < -\widetilde{v}_0 \ll -v_0^* \ll -\widetilde{v}_1 \ll -v_1^* \ll \cdots \ll -\widetilde{v}_{\frac{n-1}{2}} \ll -v^*_{\frac{n-1}{2}}$$

$$\ll -\widetilde{v}_{\frac{n-1}{2}+1} \ll \widetilde{v}_{\frac{n-1}{2}+2} \ll \cdots \ll (-1)^{\frac{n+1}{2}} \widetilde{v}_n \ll 1.$$

Note that in (3.21)

$$v^*_{\frac{n-1}{2}+i} = O(|v_0^*, v_1^*, \cdots, v^*_{\frac{n-1}{2}}|), \ i \geq 1.$$

Then, by (3.20) $\bar{M}(h)$ has $n + \frac{n+1}{2}$ positive simple zeros near $h = 0$. Hence, system (1.1) can have $n + \frac{n+1}{2}$ $(= n + [\frac{n+1}{2}]$ with n odd) limit cycles near the homoclinic loop L_0.

If n is even, by using the same method as above, we get the rank of the following matrix

$$\frac{\partial(\widetilde{v}_0, \widetilde{v}_1, \cdots, \widetilde{v}_{\frac{n-2}{2}}, \widetilde{v}_{\frac{n}{2}}, v_0^*, v_1^*, \cdots, v^*_{\frac{n-2}{2}}, \widetilde{v}_{\frac{n}{2}+1}, \widetilde{v}_{\frac{n}{2}+2}, \cdots, \widetilde{v}_n)}{\partial(a_0^+, a_2^+, \cdots, a_n^+, p_{00}^+, p_{02}^+, \cdots, p_{0,n-2}^+, a_1^-, a_3^-, \cdots, a_{n-1}^-, a_0^-, a_2^-, \cdots, a_n^-)}$$

is $n + 1 + \frac{n}{2}$, which means this matrix has a full rank in row. Thus, we can choose

$$0 < \widetilde{v}_0 \ll v_0^* \ll \widetilde{v}_1 \ll v_1^* \ll \cdots \ll \widetilde{v}_{\frac{n-2}{2}} \ll v^*_{\frac{n-2}{2}}$$

$$\ll \widetilde{v}_{\frac{n}{2}} \ll -\widetilde{v}_{\frac{n}{2}+1} \ll \cdots \ll (-1)^{\frac{n}{2}} \widetilde{v}_n \ll 1,$$

or

$$0 < -\widetilde{v}_0 \ll -v_0^* \ll -\widetilde{v}_1 \ll -v_1^* \ll \cdots \ll -\widetilde{v}_{\frac{n-2}{2}} \ll -v^*_{\frac{n-2}{2}}$$

$$\ll -\widetilde{v}_{\frac{n}{2}} \ll \widetilde{v}_{\frac{n}{2}+1} \ll \cdots \ll (-1)^{\frac{n}{2}+1} \widetilde{v}_n \ll 1.$$

Also by (3.20) and $v^*_{\frac{n-2}{2}+i} = O(|v^*_0, v^*_1, \cdots, v^*_{\frac{n-2}{2}}|),\ i \geq 1$, the function $\bar{M}(h)$ has $n + \frac{n}{2}$ $(= n + [\frac{n+1}{2}]$ with n even) positive simple zeros satisfying $0 < h \ll 1$. Therefore, system (1.9) can also have $n + [\frac{n+1}{2}]$ limit cycles near the homoclinic loop L_0 in this case. That is, $N_{homoc}(n) \geq n + [\frac{n+1}{2}]$. The proof ends.

Proof of Theorem 1.2. By Theorem 1.1, it suffices to prove that $Z(n) \leq n + [\frac{n+1}{2}]$ for $n = 1, 2, 3$ and 4.

Let $n = 1$. Then by Lemma 2.1 and (2.15)

$$\bar{M}(h) = A\sqrt{1-h} + B(1-h) + Ch\varphi_0\Big(\sqrt{\frac{1-h}{h}}\Big).$$

where A, B, C are constants. We have

$$\bar{M}(h) = \sqrt{1-h}[A + B\lambda + C\varphi_1(\lambda)],$$

where $\lambda = \sqrt{1-h} \in (0,1)$, and $\varphi_1(\lambda) = \frac{1-\lambda^2}{\lambda}\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big)$. Note that

$$\Big(\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big)\Big)' = \frac{1}{(1-\lambda^2)^2}.$$

It is direct that

$$[\varphi_1(\lambda)]' = -(1+\lambda^{-2})\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) + \frac{1}{\lambda(1-\lambda^2)},$$

and

$$[\varphi_1(\lambda)]'' = \frac{2}{\lambda^3}[\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) - \frac{\lambda}{1-\lambda^2}].$$

Note that $\frac{\lambda}{1-\lambda^2} = u\sqrt{1+u^2}$ if $u = \frac{\lambda}{\sqrt{1-\lambda^2}}$. We have $\varphi_0(u) < u\sqrt{1+u^2}$ since $(\varphi_0(u) - u\sqrt{1+u^2})' < 0$ for $u > 0$. Therefore $[\varphi_1(\lambda)]'' < 0$, which implies that $\bar{M}(h)$ has at most two simple zeros in the interval $(0,1)$. That is, $Z(1) \leq 2$.

For the case of $n = 2$, it follows from Lemma 2.1 and (2.15) that

$$\bar{M}(h) = \sqrt{1-h}[A_1 + A_2\lambda + A_3\lambda^2 + A_4\varphi_1(\lambda)],$$

where $\lambda = \sqrt{1-h} \in (0,1)$, A_1, A_2, A_3 and A_4 are constants. By simple computation, we get

$$\big[\varphi_1(\lambda)\big]''' = -\frac{6}{\lambda^4}\Big[\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) + \frac{\lambda(4\lambda^2-3)}{3(1-\lambda^2)^2}\Big] < 0,$$

since

$$\Big[\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) + \frac{\lambda(4\lambda^2-3)}{3(1-\lambda^2)^2}\Big]' = \frac{1}{(1-\lambda^2)^2} > 0.$$

Hence, $\bar{M}(h)$ has at most three simple zeros in the interval (0,1). This means $Z(2) \leq 3$.

For $n = 3$, we have

$$\bar{M}(h) = \sqrt{1-h}\Big[A_0 + A_1\lambda + A_2\lambda^2 + A_3\lambda^3 + (B_0 + B_1\lambda^2)\varphi_1(\lambda)\Big] \equiv \sqrt{1-h}M_1(\lambda).$$

Hence,

$$\begin{aligned} M_1^{(4)}(\lambda) &= 12B_1\varphi_1''(\lambda) + 8B_1\lambda\varphi_1'''(\lambda) + (B_0 + B_1\lambda^2)\varphi_1^{(4)}(\lambda) \\ &= B_1\big(12\varphi_1''(\lambda) + 8\lambda\varphi_1'''(\lambda) + \lambda^2\varphi_1^{(4)}(\lambda)\big) + B_0\varphi_1^{(4)}(\lambda). \end{aligned}$$

We can find

$$\varphi_1^{(4)}(\lambda) = \frac{24}{\lambda^5}\Big[\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) - \frac{\lambda(5\lambda^4 - 7\lambda^2 + 3)}{3(1-\lambda^2)^3}\Big],$$

and

$$12\varphi_1''(\lambda) + 8\lambda\varphi_1'''(\lambda) + \lambda^2\varphi_1^{(4)}(\lambda) = -\frac{8}{(1-\lambda^2)^3}.$$

Thus,

$$M_1^{(4)}(\lambda) = \frac{1}{(1-\lambda^2)^3}[-8B_1 + B_0(1-\lambda^2)^3\varphi_1^{(4)}(\lambda)] \equiv \frac{1}{(1-\lambda^2)^3}M_2(\lambda).$$

Since

$$\big((1-\lambda^2)^3\varphi_1^{(4)}(\lambda)\big)' = -\frac{24(1-\lambda^2)^2(\lambda^2+5)}{\lambda^6}\Big[\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) + \frac{\lambda(17\lambda^2 - 15)}{3(1-\lambda^2)^2(\lambda^2+5)}\Big]$$

and

$$\Big[\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) + \frac{\lambda(17\lambda^2 - 15)}{3(1-\lambda^2)^2(\lambda^2+5)}\Big]' = \frac{16\lambda^6}{(1-\lambda^2)^3(5+\lambda^2)^2} > 0, \tag{3.25}$$

we have for $\lambda \in (0,1)$

$$M_2'(\lambda) \neq 0 \text{ if } B_0 \neq 0.$$

Therefore, $\bar{M}(h)$ has at most 5 zeros in the interval (0,1) for this case. It follows that $Z(3) \leq 5$.

For $n = 4$, by using the same method above, we have

$$\begin{aligned} \bar{M}(h) &= \sqrt{1-h}\Big[A_0 + A_1\lambda + A_2\lambda^2 + A_3\lambda^3 + A_4\lambda^4 + (B_0 + B_1\lambda^2)\varphi_1(\lambda)\Big] \\ &\equiv \sqrt{1-h}M_3(\lambda), \end{aligned} \tag{3.26}$$

and

$$\begin{aligned} M_3^{(5)}(\lambda) &= 20B_1\varphi_1'''(\lambda) + 10B_1\lambda\varphi_1^{(4)}(\lambda) + (B_0 + B_1\lambda^2)\varphi_1^{(5)}(\lambda) \\ &= B_1\big(20\varphi_1'''(\lambda) + 10\lambda\varphi_1^{(4)}(\lambda) + \lambda^2\varphi_1^{(5)}(\lambda)\big) + B_0\varphi_1^{(5)}(\lambda), \end{aligned}$$

where

$$\varphi_1^{(5)}(\lambda) = -\frac{120}{\lambda^6}\Big[\varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) + \frac{\lambda(30\lambda^6 - 59\lambda^4 + 50\lambda^2 - 15)}{15(1-\lambda^2)^4}\Big],$$

and

$$20\varphi_1'''(\lambda) + 10\lambda\varphi_1^{(4)}(\lambda) + \lambda^2\varphi_1^{(5)}(\lambda) = -\frac{48\lambda}{(1-\lambda^2)^4}.$$

Thus,

$$M_3^{(5)}(\lambda) = \frac{\lambda}{(1-\lambda^2)^4}\Big[-48B_1 + B_0\frac{(1-\lambda^2)^4}{\lambda}\varphi_1^{(5)}(\lambda)\Big]. \tag{3.27}$$

Noting that

$$\Big[\frac{(1-\lambda^2)^4}{\lambda}\varphi_1^{(5)}(\lambda)\Big]' = -\frac{960(7-\lambda^2)(1-\lambda^2)^2}{\lambda^9}M_4(\lambda),$$

where

$$M_4(\lambda) = \varphi_0\Big(\frac{\lambda}{\sqrt{1-\lambda^2}}\Big) + \frac{\lambda(48\lambda^4-155\lambda^2+105)}{15(\lambda^2-7)(1-\lambda^2)^2},$$

we have

$$\Big[\frac{(1-\lambda^2)^4}{\lambda}\varphi_1^{(5)}(\lambda)\Big]' < 0 \tag{3.28}$$

since

$$M_4(0)=0, \quad M_4'(\lambda) = \frac{16\lambda^8(1-\lambda^2)}{5(1-\lambda^2)^4(7-\lambda^2)^2} > 0.$$

Therefore, by (3.26), (3.27) and (3.28), $\bar{M}(h)$ has at most 6 simple zeros in the interval (0,1), which yields $Z(4) \leq 6$. The proof ends.

Proof of Theorem 1.3. Following Theorem 1.1, we only need to prove $Z(n) \leq 2n + [\frac{n+1}{2}],\ n \geq 5$. We give the proof using the same method as in [11]. Let $\lambda = \sqrt{1-h},\ h \in (0,1)$. Then by (2.15) and Lemma 2.1, it follows that

$$\begin{aligned}\bar{M}(h) &= \sqrt{1-h}f_n(\sqrt{1-h}) + g_{[\frac{n-1}{2}]}(1-h)I_{10}(h)\\ &= \lambda f_n(\lambda) + (1-\lambda^2)g_{[\frac{n-1}{2}]}(\lambda^2)\varphi_0\big(\tfrac{\lambda}{\sqrt{1-\lambda^2}}\big)\\ &= \lambda f_n(\lambda) + u(\lambda)\varphi_0\big(\tfrac{\lambda}{\sqrt{1-\lambda^2}}\big)\\ &\equiv M^*(\lambda),\end{aligned}$$

where $u(\lambda) = (1-\lambda^2)u_0,\ u_0 = g_{[\frac{n-1}{2}]}(\lambda^2)$. Hence,

$$F \equiv \frac{d}{d\lambda}\big(\frac{M^*(\lambda)}{u}\big) = \Big(\frac{\lambda f_n(\lambda)}{u} + \varphi_0\big(\frac{\lambda}{\sqrt{1-\lambda^2}}\big)\Big)' = \frac{\vartheta(\lambda)}{u_0^2(1-\lambda^2)^2},$$

with

$$\vartheta(\lambda) = (\lambda f_n)'u - \lambda f_n u' + u_0^2. \tag{3.29}$$

By (3.29) and degree(u)=$2[\frac{n+1}{2}]$, we find that the coefficient of the term of $\vartheta(\lambda)$ of degree $n+2[\frac{n+1}{2}]$ vanishes if n is odd. Thus, degree(ϑ)$\leq n+2[\frac{n+1}{2}]-1$ if n is odd, and degree(ϑ)$\leq n+2[\frac{n+1}{2}]$ if n is even. That is, degree(ϑ)$\leq 2n$. Let $I=(0,1)$. Using notation $\sharp\{\lambda \in I | f(\lambda)=0\}$ to indicate the number of zeros of the function f in the interval I taking into account their multiplicities, we have

$$\sharp\{\lambda \in I | u(\lambda)=0\} \leq [\frac{n-1}{2}], \quad \sharp\{\lambda \in I|\vartheta(\lambda)=0\} \leq 2n.$$

Therefore,

$$\begin{aligned}\sharp\{\lambda\in I|M^*(\lambda)=0\} &\le \sharp\{\lambda\in I|u=0\}+\sharp\{\lambda\in I|F=0,\ u\neq 0\}+1\\ &\le \sharp\{\lambda\in I|u=0\}+\sharp\{\lambda\in I|\vartheta(\lambda)=0\}+1\\ &\le 2n+[\frac{n+1}{2}].\end{aligned}$$

The proof is completed.

Appendix

The proof of Rank($\tilde{A}_1$) $=\frac{n+1}{2}$ (for n odd). For simplicity, we take $n=7$ for example, since the method used here can be generalized for any $n\ge 1$. Thus, by (3.15) and (3.24),

$$\tilde{A}_1=\begin{pmatrix} -5!!/2^4 & -3!!/2^3 & -1/2^2 & -1/2\\ -7!!/2^5 & -5!!/2^4 & -3!!/2^3 & -1/2^2\\ -9!!/2^6 & -7!!/2^5 & -5!!/2^4 & -3!!/2^3\\ -11!!/2^7 & -9!!/2^6 & -7!!/2^5 & -5!!/2^4 \end{pmatrix}.$$

Let $r_i(c_i)$ denote the ith row(column) of $\tilde{A}_1$ for $i=1,2,3,4$. Now we making elementary transformations to $\tilde{A}_1$. Multiplying r_i by -2^i for $i=1,2,3,4$, and c_j by 2^{4-j} for $j=1,2,3$, gives

$$\tilde{A}_1\sim\bar{A}_1\equiv\begin{pmatrix} 5!! & 3!! & 1 & 1\\ 7!! & 5!! & 3!! & 1\\ 9!! & 7!! & 5!! & 3!!\\ 11!! & 9!! & 7!! & 5!! \end{pmatrix}.$$

Further, for $\bar{A}_1$, by adding r_i times $-(2i-1)$ to r_{i+1} for $i=3,2,1$, and dividing c_j by $2(4-j)$ for $j=1,2,3$, we have

$$\bar{A}_1\sim\begin{pmatrix} * & * & * & 1\\ 5!! & 3!! & 1 & 0\\ 7!! & 5!! & 3!! & 0\\ 9!! & 7!! & 5!! & 0 \end{pmatrix}.$$

Using the same method as above, we also have

$$\begin{pmatrix} * & * & * & 1\\ 5!! & 3!! & 1 & 0\\ 7!! & 5!! & 3!! & 0\\ 9!! & 7!! & 5!! & 0 \end{pmatrix}\sim\begin{pmatrix} * & * & * & 1\\ * & * & 1 & 0\\ 5!! & 3!! & 0 & 0\\ 7!! & 5!! & 0 & 0 \end{pmatrix}\sim\begin{pmatrix} * & * & * & 1\\ * & * & 1 & 0\\ * & 1 & 0 & 0\\ 5!! & 0 & 0 & 0 \end{pmatrix}.$$

Therefore, Rank($\tilde{A}_1$) $=4$. The proof ends.

References

[1] A.F. Filippov. Differential equations with discontinuous righthand sides. Kluwer Academic. Netherlands, 1988.

[2] A.A. Andronov, S.E. Khaikin, A.A. Vitt. Theory of Oscillators. Pergamon Press, Oxford, 1965.

[3] M. Kunze. Non-smooth Dynamical Systems. Spinger-Verlag, Berlin, 2000.

[4] B. Coll, A. Gasull, R. Prohens. Degenerate Hopf bifurcations in discontinuous planar systems. J. Math. Anal. Appl., 253 (2001) 671-690.

[5] A. Gasull, J. Torregrosa. Center-focus problem for discontinuous planar differential equations. Int. J. Bifur. Chaos, 13 (2003) 1755-1765.

[6] X. Chen, Z. Du. Limit cycles bifurcate from centers of discontinuous quadratic systems. Comput. Math. Appl., 59 (2010) 3836-3848.

[7] X. Liu, M. Han. Bifurcation of limit cycles by perturbing piecewise Hamiltonian systems. Internat. J. Bifur. Chaos Appl. Sci. Engrg., 5 (2010) 1-12.

[8] Z. Du, W. Zhang. Melnikov method for homoclinic bifurcation in nonlinear impact oscillators. Comput. Math. Appl., 50 (2005) 445-458.

[9] F. Battelli, M. Fečkan. Bifurcation and chaos near sliding homoclinics. J. Differential Equations, 248 (2010) 2227-2262.

[10] J. Llibre, A. Makhlonf. Bifurcation of limit cycles from a two-dimensional center inside R^n. Nonlinear Analysis, 72 (2010) 1387-1392.

[11] J. Llibre, H. Wu, J. Yu. Linear estimate for the number of limit cycles of a perturbed cubic polynomial differential system. Nonlinear Analysis, 70 (2009) 419-432.

[12] M. Han. On Hopf cyclicity of planar systems. J. Math. Anal. Appl., 245 (2000) 404-422.

[13] M. Han, G. Chen, C. Sun. On the number of limit cycles in near-Hamiltonian polynomial systems. Internat. J. Bifur. Chaos Appl. Sci. Engrg., 6 (2007) 2033-2047.

[14] Y. Wu, Y. Cao, M. Han. Bifurcations of limit cycles in a z_3-equivariant quartic planar vector field. Chaos, Solitons & Fractals, 38 (2008) 1177-1186.

[15] J. Yang, M. Han. Limit cycle bifurcations of some Liénard systems with a cuspidal loop and a homoclinic loop. Chaos, Solitons & Fractals, 44 (2011) 269-289.

[16] M. Han, W. Zhang. On Hopf bifurcation in nonsmooth planar systems. J. Differential Equations, 248 (2010) 2399-2416.

[17] M. Han, J. Chen. On the number of limit cycles in double homoclinic bifurcations. Sci. China, Ser A (2000)914-928.

[18] M. Han. Cyclicity of planar homoclinic loops and quadratic integrable systems. Sci. China, Ser. A (1997)1247-1258.